

Comparing conductance quantization in quantum wires and Quantum Hall systems

Anton Yu. Alekseev^{*†‡}, Vadim V. Cheianov^{*†‡}, Jürg Fröhlich[†]

^{*} *Institute of Theoretical Physics, Uppsala University, Box 803, S-75108, Uppsala, Sweden*

[†] *Insitut für Theoretische Physik, ETH-Hönggerberg, CH-8093, Zürich, Switzerland*

[‡] *Steklov Mathematical Institute, Fontanka 27, 191011, St.Petersburg, Russia*

(July 1996)

We propose a new calculation of the DC conductance of a 1-dimensional electron system described by the Luttinger model. Our approach is based on the ideas of Landauer and Büttiker and on the methods of current algebra. We analyse in detail the way in which the system can be coupled to external reservoirs. This determines whether the conductance is renormalized or not. We show that although a quantum wire and a Fractional Quantum Hall system are described by the same effective theory, their coupling to external reservoirs is different. As a consequence, the conductance in the wire is quantized in integer units of e^2/h per spin orientation whereas the Hall conductance allows for fractional quantization.

Recent experiments on transport properties of quantum wires [1,2] attracted new interest to the problem of conductance of a 1-dimensional electron gas.

Since the work of Apel and Rice [3], where the authors computed the conductance in the 1-dimensional Luttinger model, it was believed that the effects of interaction in the one-channel quantum wire should lower the conductance with respect to the value

$$\sigma = 2 \frac{e^2}{h} \quad (1)$$

predicted by the Landauer-Büttiker formula [4] for the case of non-interacting electrons. However, experimental data suggest that this renormalization of conductance does not actually take place [2].

At the moment there exist a number of theoretical arguments [5–9] intended to explain the non-renormalization of the conductance in 1-dimensional electron gas. One of them [5,6] is based on the idea that the conductance of the 1-channel quantum wire is completely determined by the structure of the leads to which the wire is attached. So far as the electrons in the leads form a Landau Fermi-liquid one chooses to model the leads by two semi-infinite 1-dimensional 1-channel non-interacting Fermi liquids whereas the wire is modeled by a Luttinger Fermi-liquid. Due to strong non-locality of current-current correlators in 1-dimensional systems the calculation based on the Kubo formula leads to the non-renormalized result (1).

Another approach [8,9] is based on the observation that the dielectric constant in the Luttinger Fermi-liquid increases with increasing strength of interaction. This fact changes the definition of the voltage drop which results in the cancellation of the renormalization factor in the conductance. In this approach one does not consider the reservoirs to which the wire is attached and does not explain how the voltage drop is measured in experiment.

Both approaches mentioned above are based on linear response theory and its main tool, the Kubo formula, in their calculation of conductance. Yet there is another, in many ways more intuitive approach based on ideas of

Landauer and Büttiker [4]. In this paper we show how to apply these ideas to a Luttinger Fermi-liquid type channel.

We study how the Luttinger liquid interacts with external reservoirs. In order to clarify our analysis of quantum wires, we provide a parallel treatment of the Fractional Quantum Hall system which is described by the same effective Hamiltonian [10,11]. It appears that the electron transfer between the Luttinger liquid and external reservoirs in these two systems is organized in a different way. This difference leads to the integer quantization of conductances in quantum wires in contrast to the fractional quantization of Hall conductances.

We replace appealing but clearly unrealistic (1-dimensional) models of the leads [5,6] by a more universal consideration which relies only on the fact that the leads interact with the Luttinger liquid via electron transfer. It is difficult to compare our results to [8,9] as they do not consider the interaction to reservoirs at all and in our approach this is the central issue which determines whether the conductance is renormalized or not.

It is well known that a 1-dimensional interacting electron system is effectively described by the Luttinger model [12]. For simplicity we consider spinless fermions. This makes the comparison to the spin polarized Quantum Hall system more straightforward. Polarization of fermions can be easily recovered in the final expression for conductance.

The bosonized Lagrangian density of the Luttinger model looks as follows [12]:

$$\mathcal{L} = \frac{\hbar v_F}{4} \left(\frac{1}{v_F^2} (\partial_t \phi)^2 - (1 + g) (\partial_x \phi)^2 \right), \quad (2)$$

where v_F is the Fermi velocity, $n = \partial_x \phi$ is the particle density, $I = e \partial_t \phi$ is the electric current, g is an effective coupling constant. The corresponding Hamiltonian is given by

$$H_0 = \frac{\hbar v_F}{4} \int dx : \left(\frac{1}{v_F^2} (\partial_t \phi)^2 + (1 + g) (\partial_x \phi)^2 \right) :, \quad (3)$$

where $::$ stands for the standard Wick ordering. The

model with $g = 0$ describes a non-interacting 1 - dimensional Fermi-liquid.

The Hamiltonian (3) can be expressed in terms of left-moving and right-moving currents:

$$H_0 = \frac{\hbar}{2v_F} \int dx : (I_L^2 + I_R^2) : , \quad (4)$$

where

$$\begin{aligned} I_L &= \frac{1}{2}(\partial_t \phi + v_F \sqrt{1+g} \partial_x \phi), \\ I_R &= -\frac{1}{2}(\partial_t \phi - v_F \sqrt{1+g} \partial_x \phi), \end{aligned} \quad (5)$$

so that

$$I = e(I_L - I_R) , \quad n = \frac{1}{v_F \sqrt{1+g}}(I_L + I_R). \quad (6)$$

Let us remark that the same effective Hamiltonian describes edge excitations of a spin polarized incompressible Quantum Hall fluid on a cylinder [10] with one chiral channel of edge excitations per boundary. Then eI_L and eI_R are the edge currents on the two boundaries of the cylinder. The filling factor of the Quantum Hall fluid is related to the effective coupling of the Luttinger model via:

$$\nu = \frac{1}{\sqrt{1+g}}. \quad (7)$$

Incompressibility requires that ν^{-1} is an odd integer.

In order to measure conductances in the Luttinger model we should couple it to two reservoirs with different chemical potentials [4]. This can be done in two different ways. We shall see that one way is realized in the Quantum Hall system. The corresponding conductance should be interpreted as the Hall conductance

$$\sigma_H = \nu \frac{e^2}{h}. \quad (8)$$

The other way of coupling the Luttinger model to reservoirs is realized in thin wires and leads to the non-renormalized value

$$\sigma = \frac{e^2}{h}. \quad (9)$$

The coupling of reservoirs to the Luttinger liquid can only be organized via electron transfer. We therefore must identify excitations in the Luttinger liquid which correspond to electrons and assign to them chemical potentials of reservoirs according to the conditions of the measurement.

This can be done naturally for incompressible Quantum Hall fluids. In this case the chiral channels are spatially separated by the bulk of the sample and the reservoirs are attached directly to each of them. Mathematically this is expressed by adding an extra term to the Hamiltonian:

$$H_{\text{Hall}} = H_0 + \mu_L N_L + \mu_R N_R, \quad (10)$$

where μ_L and μ_R are the chemical potentials of the reservoirs $eV = \mu_R - \mu_L$, N_L and N_R are conserved electron numbers in the left-moving and right-moving channels (corresponding to the two different edges of the sample):

$$\begin{aligned} N_L &= \int dx n_L , \quad n_L = \frac{1}{v_F} \nu I_L; \\ N_R &= \int dx n_R , \quad n_R = \frac{1}{v_F} \nu I_R. \end{aligned} \quad (11)$$

The expectation value of the operator (10) can be interpreted as a free energy of the system at zero temperature. We minimize the energy functional which yields the uniform ground state current

$$I = e \nu \left(\frac{\mu_R}{h} - \frac{\mu_L}{h} \right) = \nu \frac{e^2}{h} V. \quad (12)$$

Thus, the Hall conductance is given by formula (8). This result would correspond to the renormalized conductance

$$\sigma = \frac{1}{\sqrt{1+g}} \frac{e^2}{h}$$

in the Luttinger model.

In the Quantum Hall system the left- and right-moving edge excitations of charge e are represented by operators

$$\begin{aligned} \Psi_L(x) &=: \exp\{i\pi(\sqrt{1+g}\phi(x) + \frac{1}{v_F} \int^x \partial_t \phi dx)\} : , \\ \Psi_R(x) &=: \exp\{i\pi(\sqrt{1+g}\phi(x) - \frac{1}{v_F} \int^x \partial_t \phi dx)\} : . \end{aligned} \quad (13)$$

These operators correspond to particles with Fermi statistics only when $\sqrt{1+g}$ is an odd integer [10,11]. These values of the filling factor

$$\nu = \frac{1}{2m+1} \quad (14)$$

correspond to Laughlin's plateaux [13] in the theory of Fractional Quantum Hall Effect.

Next we turn to the analysis of quantum wires. For generic values of the coupling constant g neither left- nor right-moving edge excitations can be identified with electrons. Hence, we can not assign particular values of chemical potentials to these excitation branches. Yet, there always exist other fermionic excitations of charge e that describe the physical electrons in the Luttinger model. They are created by applying the operators Ψ_+ , Ψ_- given by

$$\begin{aligned} \Psi_+(x) &=: \exp\{i\pi(\phi(x) + \frac{1}{v_F} \int^x \partial_t \phi dx)\} : , \\ \Psi_-(x) &=: \exp\{i\pi(\phi(x) - \frac{1}{v_F} \int^x \partial_t \phi dx)\} : . \end{aligned} \quad (15)$$

These operators coincide with Ψ_L , Ψ_R in the non-interacting model ($g = 0$). The particle densities corresponding to the operators Ψ_+ and Ψ_- are given by

$$\begin{aligned} n_+ &= \frac{1 + \sqrt{1+g}}{2} n_L + \frac{1 - \sqrt{1+g}}{2} n_R, \\ n_- &= \frac{1 - \sqrt{1+g}}{2} n_L + \frac{1 + \sqrt{1+g}}{2} n_R. \end{aligned} \quad (16)$$

It is natural to view the chemical potentials of the reservoirs as the variables conjugate to the conserved charges

$$N_+ = \int dx n_+, \quad N_- = \int dx n_- \quad (17)$$

which measure the total number of electrons created by powers of Ψ_+ and Ψ_- , respectively. If the quantum channel in the wire is ballistic and adiabatic, the chemical potential of the left reservoir is conjugate to N_+ whereas the chemical potential of the right reservoir is conjugate to N_- . The counterpart of the Hamiltonian (10) therefore is given by

$$H_{\text{Wire}} = H_0 + \mu_L N_+ + \mu_R N_- \quad (18)$$

A calculation similar to the one in (12) shows that the ground state of H_{Wire} carries a current

$$I = e \left(\frac{\mu_R}{h} - \frac{\mu_L}{h} \right) = \frac{e^2}{h} V \quad (19)$$

corresponding to (9). Two spin polarizations of electrons can be included in the final answer for conductance of a single quantum channel by adding a factor of 2 which leads to (1).

We conclude that although the Quantum Hall system and the quantum wire are described by the same effective Hamiltonian, the way in which they are coupled to reservoirs in transport measurements is different. Mathematically, this is reflected in using chemical potentials $\mu_L(\mu_R)$ of left(right) reservoirs conjugate variables to different conserved charges $N_L(N_R)$, for Quantum Hall fluids, and $N_+(N_-)$ for quantum wires. As a consequence, in the Quantum Hall system the conductance depends on the effective coupling constant related to the filling factor by (7), but this is not the case for the quantum wire.

-
- [1] B.J. van Wees *et al.*, Phys. Rev. Lett. **60**, 848 (1988).
 - [2] S.Tarucha, T.Honda, and T.Saku, Solid State Commun. **94**, 413 (1995).
 - [3] W.Apel and T.M.Rice, Phys. Rev. B **26**, 7063 (1982).
 - [4] R.Landauer, Philos. Mag. **21** 863 (1970);
M.Büttiker, Phys. Rev. Lett. **57**, 1761 (1986).
 - [5] V.V.Ponomarenko, Phys. Rev. B **52**, R8666 (1995).
 - [6] D.L.Maslov and M.Stone, Phys. Rev. B **52**, R5539 (1995).
 - [7] I.Safi and H.J.Schulz, Phys. Rev. B **52**, R17040 (1995).
 - [8] A.Kawabata, J. Phys. Soc. Japan **65**, 30 (1996).
 - [9] S.Fujimoto and N.Kawakami, con-mat/9602101.
 - [10] X.G.Wen, Phys. Rev. B **41**, 12838 (1990).
 - [11] J.Fröhlich and T.Kerler, Nucl. Phys. B **354**, 369 (1991).
J.Fröhlich, T.Kerler, U.Studer, E.Thiran, Nucl. Phys. B **453** [FS], 670 (1995).
 - [12] J.Sólyom, Adv. Phys. **28**, 209 (1979).
F.D.M.Haldane, J. Phys. C **14**, 2585 (1981).
 - [13] R.B.Laughlin, Phys. Rev. Lett. **50**, 1395 (1983); Phys. Rev. B **27**, 3383 (1983).